

N-Level Math (4045) Formula List

*Formulas highlighted in yellow are found in the formula list of the exam paper.

Unit Conversion	
<p>Area</p> $1\text{m}^2 = 100\text{cm} \times 100\text{cm}$ $= 10\,000\text{cm}^2$ <p>Volume</p> $1\text{m}^3 = 100\text{cm} \times 100\text{cm} \times 100\text{cm}$ $= 1\,000\,000\text{cm}^3$ <p>1 Litre = 1000cm^3 (since $1\text{mg} = 1\text{cm}^3$)</p>	$1\text{km}^2 = 1000\text{m} \times 1000\text{m}$ $= 1\,000\,000\text{m}^2$ <div style="text-align: center;"> <p>km/h $\xrightarrow{\times \frac{10}{36}}$ m/s m/s $\xrightarrow{\times \frac{36}{10}}$ km/h</p> </div>

Financial Math		
<p>Percentage Increase</p> $= \frac{\text{Increase}}{\text{Original}} \times 100\%$	<p>Percentage Decrease</p> $= \frac{\text{Decrease}}{\text{Original}} \times 100\%$	
<p>Simple Interest</p> $I = \frac{P \times R \times T}{100}$	<p>Compound Interest</p> $P + I = P \times \left(1 + \frac{R}{100}\right)^n$	<p>P-Principal (\$)</p> <p>I- Interest Amount (\$)</p> <p>R-Interest Rate (%)</p> <p>T-Time (Years)</p> <p>n-No. of Periods</p>

Ratio and Proportion

Conversion between Length (Linear) and Area Ratio.

$$\frac{(L_s)^2}{(L_b)^2} = \frac{A_s}{A_b}$$

$$\frac{\sqrt{A_s}}{\sqrt{A_b}} = \frac{L_s}{L_b}$$

Conversion between Length (Linear) and Volume Ratio.

$$\frac{(L_s)^3}{(L_b)^3} = \frac{V_s}{V_b}$$

$$\frac{\sqrt[3]{V_s}}{\sqrt[3]{V_b}} = \frac{L_s}{L_b}$$

L_s : Length-smaller object:

L_b : Length-bigger object

A_s : Area-smaller object

A_b : Area-bigger object

V_s : Volume-smaller object

V_b : Volume-bigger object

To convert Area to Volume & vice versa, first convert to Length.

$$\left(\frac{A_s}{A_b} \right) \begin{array}{c} \xrightarrow{\sqrt{\text{Square Root}}} \\ \xleftarrow{\text{Square}^2} \end{array} \left(\frac{L_s}{L_b} \right) \begin{array}{c} \xrightarrow{\text{Cube}^3} \\ \xleftarrow{\sqrt[3]{\text{Cube Root}}} \end{array} \left(\frac{V_s}{V_b} \right)$$

Speed and Distance

$$\text{Speed} = \frac{\text{Distance}}{\text{Time}} \quad \text{Acceleration} = \frac{\text{Final Speed} - \text{Initial Speed}}{\text{Time Taken}}$$

$$\text{Average Speed} = \frac{\text{Total Distance}}{\text{Total Time}}$$

Distance Travelled is the Area UNDER the **speed /time** graph

N-Level Math (4045) Formula List

Indices	
$x^a \times x^b = x^{a+b}$ $a^m \times b^m = (a \times b)^m$	<p style="color: green;">Base No. same → Power add</p> <p style="color: green;">Power same → Base No. multiply</p>
$\frac{x^a}{x^b} = x^{a-b}$ $\frac{a^m}{b^m} = \left(\frac{a}{b}\right)^m$	<p style="color: green;">Base No. same → Power minus</p> <p style="color: green;">Power same → Base No. divide</p>
<p style="color: green;">Note: $(x^a)^b = x^{a \times b}$</p> <p style="color: green;">$(x^a)^b \neq x^{a+b}$</p>	$x^0 = 1$
$x^{-a} = \frac{1}{x^a} \qquad \frac{1}{x^{-a}} = x^a \qquad x^a y^{-b} = \frac{x^a}{y^b}$	
$\left(\frac{x}{y}\right)^{-a} = \left(\frac{y}{x}\right)^a \qquad x^{\frac{1}{b}} = \sqrt[b]{x^1} \qquad x^{\frac{a}{b}} = \sqrt[b]{x^a}$	
$x^{-\frac{a}{b}} = \frac{1}{x^{\frac{a}{b}}} = \frac{1}{\sqrt[b]{x^a}} \qquad x^{-\frac{1}{b}} = \frac{1}{x^{\frac{1}{b}}} = \frac{1}{\sqrt[b]{x^1}}$	


Coordinate Geometry

Linear Graph

$Y = m x + c$ where $m =$ gradient

and $c =$ y-intercept

$$\text{Gradient}(m) = \frac{y_2 - y_1}{x_2 - x_1}$$



Mid-point of a line

$$= \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Distance between two points

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Polygons

Sum of interior angles of an n-sided polygon.

$$(n - 2) \times 180^{\circ}$$

Each interior angle of an n-sided polygon.

$$\frac{(n - 2) \times 180^{\circ}}{n}$$

Each exterior angle of an n-sided polygon

$$= \frac{360^{\circ}}{n}$$

Sum of exterior angles of an n-sided polygon = 360°


Arc Length, Sector and Segment

Arc Length

$$S = \frac{\theta^{\circ}}{360^{\circ}} \times 2\pi r \text{ (Degree) Or } S = r\theta \text{ (Radian)}$$

Area of Sector

$$A = \frac{\theta^{\circ}}{360^{\circ}} \times \pi r^2 \text{ (Degree) Or } A = \frac{1}{2} r^2 \theta \text{ (Radian)}$$




Mensuration

Circles

$$\text{Area} = \pi \times r^2$$

$$\text{Circumference} = 2 \times \pi \times r$$


$$\text{Or } \pi \times d$$



Trapezium

$$\text{Perimeter} = w + x + y + z$$


$$\text{Area} = \frac{1}{2} \times (x + y) \times h$$



Parallelogram

$$\text{Perimeter} = 2 \times y + 2 \times z$$

$$\text{Area} = y \times z$$



Cylinder

Total Surface Area

(close cylinder)


$$= 2 \times \pi \times r^2 + 2 \times \pi \times r \times h$$

Total Surface Area

(open cylinder)

$$= \pi \times r^2 + 2 \times \pi \times r \times h$$

$$\text{Volume} = \pi \times r^2 \times h$$



Cone

Total Surface Area =


$$\pi \times r \times l + \pi \times r^2$$

$$\text{Volume} = \frac{1}{3} \times \pi \times r^2 \times h$$

l=slant height

h=vertical height

(Note the difference)



Pyramid


Total Surface Area =

Sum of 4 triangles + base

$$\text{Volume} = \frac{1}{3} \times A \times h$$

A=base area

h=vertical height



Sphere


$$\text{Total Surface Area} = 4 \times \pi \times r^2$$

$$\text{Volume} = \frac{4}{3} \times \pi \times r^3$$

Hemisphere (half-sphere)





$$\text{Total Surface Area} = 2 \times \pi \times r^2 + \pi \times r^2$$

$$\text{Volume} = \frac{2}{3} \times \pi \times r^3$$




Properties of Circle

Angle at Centre = Twice Angle at Circumference




Angles in the Same Segment

(Are Equal)



Angle in a Semi-circle = 90°




Angles in Opposite Segment

(Add up to 180°)


$$a^\circ + c^\circ = 180^\circ$$

$$b^\circ + d^\circ = 180^\circ$$




Exterior angle of a cyclic quadrilateral

$$b^\circ = p^\circ$$




Isosceles Triangle




**Perpendicular from Centre
Bisects Chord**

$$\angle OXA = \angle OXB = 90^\circ$$




**Equal Chord, Equal Distance
from Centre**



Tangent Perpendicular Radius

$$\angle OAB = 90^\circ$$




Tangents from External Point

$$BC = BA$$

$$\angle OCB = \angle OAB = 90^\circ$$

$$OA = OC \text{ (radius)}$$




Trigonometry

Note: Use when the triangle is Right Angle.

$$\tan B = \frac{\text{Opp (DE)}}{\text{Adj (EB)}} \text{ (TOA)}$$

$$\cos B = \frac{\text{Adj (EB)}}{\text{Hyp (DB)}} \text{ (CAH)}$$

$$\sin B = \frac{\text{Opp (DE)}}{\text{Hyp (DB)}} \text{ (SOH)}$$



Pythagoras Theorem $DB^2 = DE^2 + EB^2$

Note: Use when the triangle is NOT Right Angle.


$$\text{Area of Triangle} = \frac{1}{2} \times a \times b \times \sin C$$

Sine Rule

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \times \cos C$$



Probability

$$\text{Probability} = \frac{\text{Number Of Successful Outcome}}{\text{Total Number Of Outcomes}} \quad \text{OR} \quad \frac{\text{Success}}{\text{Success} + \text{Failure}}$$

If the probability of A **AND** B occurs, then $P(A) \times P(B)$.

If the probability of A **OR** B occurs, then $P(A) + P(B)$

If the probability of A **DOES NOT** occurring, then $1 - P(A)$.

Probability is between and include 0 to 1.

If Probability $(P) = 0$, it means that there is **NO CHANCE** of success.

If Probability $(P) = 1$ it means that success is **CERTAIN**.

Statistics

Ungroup Data

$$\text{Mean}(\bar{X}) = \frac{\text{Sum Of All Data Values}}{\text{Number Of Data}}$$

Group Data

$$\text{Mean}(\bar{X}) = \frac{\sum fx}{\sum f}$$

$$\text{Lower Quartile} = \frac{1}{4}(n + 1)\text{th Term}$$

n is the total frequency.

N-Level Math (4045) Formula List

<p>$Median = \left(\frac{n+1}{2}\right)th \text{ Term}$</p> <p>$Upper \text{ Quartile} = \frac{3}{4}(n+1)th \text{ Term}$</p>	<p>*These formulas give the POSITION of the value in ascending order. It DOES NOT give the actual value.</p>
<p>Ungroup Data - Standard Deviation (σ)</p> $\sigma = \sqrt{\frac{\sum (x - \bar{X})^2}{\sum f}} \quad \text{or} \quad \sigma = \sqrt{\frac{\sum x^2}{n} - \bar{X}^2}$	
<p>Group Data - Standard Deviation (σ)</p> $\sigma = \sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2} \quad \text{or} \quad \sigma = \sqrt{\frac{\sum fx^2}{\sum f} - (\bar{X})^2}$	
<p>The End</p>	